

5-1-2010

Two-loop Resummations for QCD Hard Scattering

Nikolaos Kidonakis

Kennesaw State University, nkidonak@kennesaw.edu

Follow this and additional works at: <https://digitalcommons.kennesaw.edu/facpubs>



Part of the [Physics Commons](#)

Recommended Citation

Kidonakis, Nikolaos, "Two-loop Resummations for QCD Hard Scattering" (2010). *Faculty Publications*. 3873.
<https://digitalcommons.kennesaw.edu/facpubs/3873>

This Article is brought to you for free and open access by DigitalCommons@Kennesaw State University. It has been accepted for inclusion in Faculty Publications by an authorized administrator of DigitalCommons@Kennesaw State University. For more information, please contact digitalcommons@kennesaw.edu.

Two-loop resummation for QCD hard scattering

Nikolaos Kidonakis^{*†}

Kennesaw State University, USA

E-mail: nkidonak@kennesaw.edu

I present results for the resummation of soft and collinear gluon contributions to QCD hard-scattering cross sections at two loops. This requires the calculation of UV and IR divergences through two loops and the construction of soft anomalous dimensions that control resummation at NNLL. Explicit results are provided for processes that involve massless and/or massive quarks, including single top and top pair production.

XVIII International Workshop on Deep-Inelastic Scattering and Related Subjects

April 19 -23, 2010

Convitto della Calza, Firenze, Italy

^{*}Speaker.

[†]This work was supported by the National Science Foundation under Grant No. PHY 0855421.




Figure 1: Two-loop vertex-correction graphs.

1. Resummation

Soft-gluon corrections are important in many QCD processes, particularly near threshold. The calculation of these corrections is needed at higher orders for increased accuracy in theoretical predictions. Terms of the form $[\ln^k(s_4/M^2)/s_4]_+$, where $s_4 \rightarrow 0$ at threshold, arise from incomplete cancellations of infrared divergences between virtual diagrams and real diagrams with soft (low-energy) gluons. These soft-gluon corrections exponentiate. The resummation follows from factorization properties of the cross section. At next-to-leading-logarithm (NLL) accuracy resummation requires one-loop calculations in the eikonal approximation; at next-to-next-to-leading-logarithm (NNLL) accuracy it requires two-loop calculations.

The resummation is performed in moment space, with N the moment variable conjugate to s_4 . We use renormalization group evolution (RGE) to evolve the soft-gluon function associated with soft-gluon emission. The resummed cross section is

$$\hat{\sigma}^{res}(N) = \exp[E(N)] \operatorname{tr} \left\{ H(\sqrt{s}) \exp \left[\int_{\sqrt{s}}^{\sqrt{s}/\tilde{N}} \frac{d\mu}{\mu} \Gamma_S^\dagger(\mu) \right] S \left(\frac{\sqrt{s}}{\tilde{N}} \right) \exp \left[\int_{\sqrt{s}}^{\sqrt{s}/\tilde{N}} \frac{d\mu}{\mu} \Gamma_S(\mu) \right] \right\} \quad (1.1)$$

where the exponent E resums universal collinear and soft gluon emission [1, 2] from the incoming and outgoing partons, H is the N -independent hard-scattering function, and S is the soft-gluon function whose RGE is governed by Γ_S , the soft anomalous dimension [3, 4]. The functions H , S , and Γ_S are matrices in color-exchange space.

We calculate Γ_S in the eikonal approximation, in which the Feynman rules for the quark-gluon vertex give $g_s T_F^c (v^\mu / v \cdot k)$, with $g_s^2 = 4\pi\alpha_s$, v a dimensionless velocity vector, k the gluon momentum, and T_F^c the generators of $SU(3)$. We perform the calculations in momentum space and Feynman gauge. We expand the soft anomalous dimension as $\Gamma_S = (\alpha_s/\pi)\Gamma_S^{(1)} + (\alpha_s/\pi)^2\Gamma_S^{(2)} + \dots$, and determine it through two loops for a number of processes.

2. Massive soft (cusp) anomalous dimension

We begin with two-loop results for the soft (cusp [5]) anomalous dimension for $e^+e^- \rightarrow t\bar{t}$

[6]. Calculations with massive quarks present considerable complications in addition to those with massless quarks [7]. The two-loop vertex-correction eikonal diagrams are shown in Fig. 1; there are additional diagrams with top-quark self energies [6]. We include counterterms for all graphs and multiply with corresponding color factors. Ultraviolet (UV) and infrared (IR) singularities are separated, and we determine the two-loop soft anomalous dimension from the UV poles of the sum of the graphs. In terms of the cusp angle [5] $\gamma = \ln[(1 + \beta)/(1 - \beta)]$, with $\beta = \sqrt{1 - 4m^2/s}$ and m the heavy quark mass, we get $\Gamma_S^{(1)} = C_F(\gamma \coth \gamma - 1)$ and [6]

$$\Gamma_S^{(2)} = \frac{K}{2}\Gamma_S^{(1)} + C_F C_A \left\{ \frac{1}{2} + \frac{\zeta_2}{2} + \frac{\gamma^2}{2} - \frac{1}{2} \coth^2 \gamma \left[\zeta_3 - \zeta_2 \gamma - \frac{\gamma^3}{3} - \gamma \text{Li}_2(e^{-2\gamma}) - \text{Li}_3(e^{-2\gamma}) \right] \right. \\ \left. - \frac{1}{2} \coth \gamma \left[\zeta_2 + \zeta_2 \gamma + \gamma^2 + \frac{\gamma^3}{3} + 2\gamma \ln(1 - e^{-2\gamma}) - \text{Li}_2(e^{-2\gamma}) \right] \right\} \quad (2.1)$$

where $K = C_A(67/18 - \zeta_2) - 5n_f/9$. $\Gamma_S^{(2)}$ vanishes at $\beta = 0$, the threshold limit, and diverges at $\beta = 1$, the massless limit. In the case where one quark is massive and one is massless the result simplifies and we find $\Gamma_S^{(2)} = (K/2)\Gamma_S^{(1)} + C_F C_A(1 - \zeta_3)/4$.

3. Single top quark processes

The color structure gets more complicated with more than two colored partons in a hard-scattering process. The cusp anomalous dimension, presented in the previous section, is an essential component of calculations for other processes. Next, we compute two-loop soft anomalous dimensions for single top production via the s -channel, and for associated production of a top quark with a W boson or a charged Higgs.

3.1 s -channel single top quark production

For s -channel single-top production the diagrams involve different pairings of the two-loop diagrams for the cusp anomalous dimension. We find the one-loop result for the soft anomalous dimension [8, 9]

$$\Gamma_{S, \text{top } s\text{-ch}}^{(1)} = C_F \left[\ln \left(\frac{s - m_t^2}{m_t \sqrt{s}} \right) - \frac{1}{2} \right] \quad (3.1)$$

and the two-loop [9] expression

$$\Gamma_{S, \text{top } s\text{-ch}}^{(2)} = \frac{K}{2}\Gamma_{S, \text{top } s\text{-ch}}^{(1)} + C_F C_A \frac{(1 - \zeta_3)}{4}. \quad (3.2)$$

Using this result in Eq. (1.1) and expanding the resummed cross section to next-to-next-to-leading order (NNLO) in the strong coupling, we derive an approximate NNLO cross section. As shown in [9, 10] the NNLO approximate corrections enhance the NLO cross section by about 15% at the Tevatron and about 13% at LHC energies.

3.2 Associated production of a top quark with a W^- or H^-

For the processes $bg \rightarrow tW^-$ or $bg \rightarrow tH^-$ the two-loop vertex-correction eikonal diagrams are shown in Fig. 2 (there are additional top-quark self-energy graphs not shown here). Note




Figure 2: Eikonal two-loop graphs for tW^- or tH^- production.

that graphs with gluons attached to all three eikonal lines, not shown here, either vanish or do not contribute to the two-loop soft anomalous dimension. The one-loop result for the soft anomalous dimension for $bg \rightarrow tW^-$ is

$$\Gamma_{S,tW^-}^{(1)} = C_F \left[\ln \left(\frac{m_t^2 - t}{m_t \sqrt{s}} \right) - \frac{1}{2} \right] + \frac{C_A}{2} \ln \left(\frac{m_t^2 - u}{m_t^2 - t} \right) \quad (3.3)$$

and at two loops we find

$$\Gamma_{S,tW^-}^{(2)} = \frac{K}{2} \Gamma_{S,tW^-}^{(1)} + C_F C_A \frac{(1 - \zeta_3)}{4}. \quad (3.4)$$

The NNLO approximate corrections for tW^- production enhance the NLO cross section by about 8% at LHC energies [10].

The same analytical result for Γ_S holds for $bg \rightarrow tH^-$ [10, 11]. The NNLO approximate corrections for tH^- production provide around 15% to 20% enhancement over NLO at LHC energies [10].

Related results can also be derived for direct photon production.

4. Top-antitop pair hadroproduction

Resummation for $t\bar{t}$ production at hadron colliders at NLL accuracy was presented in [3] where the one-loop soft anomalous dimension matrix was first calculated. Detailed numerical studies including the NLL terms and sets of subleading terms [12] have shown that threshold resummation increases the cross section of $t\bar{t}$ production and significantly reduces the scale dependence at both Tevatron and LHC energies.

The two-loop results of [6] are an essential ingredient in determining the complete two-loop matrix result for $t\bar{t}$ hadroproduction. If we rewrite Eq. (2.1) as $\Gamma_S^{(2)} = (K/2)\Gamma_S^{(1)} + C_F C_A M_\beta$ (i.e. M_β denotes the terms inside the curly brackets in Eq. (2.1), with the cusp angle γ expressed explicitly in terms of β), then the diagonal elements of the two-loop soft anomalous dimension matrix, for both $q\bar{q} \rightarrow t\bar{t}$ and $gg \rightarrow t\bar{t}$ channels, can be written as a multiple of the one-loop result (in Feynman gauge) plus a multiple of M_β . For example, in a singlet-octet basis for the $q\bar{q} \rightarrow t\bar{t}$ channel we have

$$\Gamma_{q\bar{q}11}^{(2)} = \frac{K}{2} \Gamma_{q\bar{q}11}^{(1)} + C_F C_A M_\beta,$$

$$\Gamma_{q\bar{q}22}^{(2)} = \frac{K}{2}\Gamma_{q\bar{q}22}^{(1)} + C_A \left(C_F - \frac{C_A}{2} \right) M_\beta. \quad (4.1)$$

Similar results are found for the gg channel. The off-diagonal elements obey a similar relation in terms of a function N_β which involves a subset of the terms in M_β . Complete results will appear in a forthcoming paper. It is important to note that the inclusion of the complete two-loop expressions for the soft anomalous dimension matrix do not change the results of [12] by more than $O(\text{per mille})$, thus showing the robustness of those earlier results and proving that they captured the dominant subleading terms as was explained in detail in [12].

References

- [1] G. Sterman, *Summation of Large Corrections to Short Distance Hadronic Cross-Sections*, *Nucl. Phys. B* **281**, 310 (1987).
- [2] S. Catani and L. Trentadue, *Resummation of the QCD Perturbative Series for Hard Processes*, *Nucl. Phys. B* **327**, 323 (1989).
- [3] N. Kidonakis and G. Sterman, *Subleading logarithms in QCD hard scattering*, *Phys. Lett. B* **387**, 867 (1996); *Resummation for QCD hard scattering*, *Nucl. Phys. B* **505**, 321 (1997) [hep-ph/9705234].
- [4] N. Kidonakis, *Next-to-next-to-next-to-leading-order soft-gluon corrections in hard-scattering processes near threshold*, *Phys. Rev. D* **73**, 034001 (2006) [hep-ph/0509079].
- [5] G.P. Korchemsky and A.V. Radyushkin, *Loop-Space Formalism And Renormalization Group For The Infrared Asymptotics Of QCD*, *Phys. Lett. B* **171**, 459 (1986); *Renormalization of the Wilson Loops Beyond the Leading Order*, *Nucl. Phys. B* **283**, 342 (1987).
- [6] N. Kidonakis, *Two-Loop Soft Anomalous Dimensions and Next-to-Next-to-Leading-Logarithm Resummation for Heavy Quark Production*, *Phys. Rev. Lett.* **102**, 232003 (2009), arXiv:0903.2561 [hep-ph]; *Two-loop soft anomalous dimensions with massive and massless quarks*, in proceedings of *DPF 2009*, eConf C090726, arXiv:0910.0473 [hep-ph].
- [7] S.M. Aybat, L.J. Dixon, and G. Sterman, *Two-loop soft anomalous dimension matrix and resummation at next-to-next-to-leading poles*, *Phys. Rev. D* **74**, 074004 (2006) [hep-ph/0607309].
- [8] N. Kidonakis, *Single top quark production at the Fermilab Tevatron: Threshold resummation and finite-order soft gluon corrections*, *Phys. Rev. D* **74**, 114012 (2006) [hep-ph/0609287]; *Higher-order soft gluon corrections in single top quark production at the CERN LHC*, *Phys. Rev. D* **75**, 071501(R) (2007) [hep-ph/0701080].
- [9] N. Kidonakis, *Next-to-next-to-leading logarithm resummation for s-channel single top quark production*, *Phys. Rev. D* **81**, 054028 (2010), arXiv:1001.5034 [hep-ph].
- [10] N. Kidonakis, *Single top quark production cross section at hadron colliders*, in these proceedings, *PO S (DIS 2010) 196*, arXiv:1005.3330 [hep-ph]; and in preparation.
- [11] N. Kidonakis, *Charged Higgs production via $b\bar{g} \rightarrow tH^-$ at the LHC*, *JHEP* **05** (2005) 011 [hep-ph/0412422].
- [12] N. Kidonakis and R. Vogt, *Next-to-next-to-leading order soft-gluon corrections in top quark hadroproduction*, *Phys. Rev. D* **68**, 114014 (2003) [hep-ph/0308222]; *Theoretical top quark cross section at the Fermilab Tevatron and the CERN LHC*, *Phys. Rev. D* **78**, 074005 (2008), arXiv:0805.3844 [hep-ph].